




ELSEVIER

Contents lists available at ScienceDirect

Chaos, Solitons & Fractals

Nonlinear Science, and Nonequilibrium and Complex Phenomena

journal homepage: www.elsevier.com/locate/chaos

Simple chaotic flows with a line equilibrium

Sajad Jafari^{a,*}, J.C. Sprott^b^a Biomedical Engineering Department, Amirkabir University of Technology, Tehran 15875-4413, Iran^b Department of Physics, University of Wisconsin, Madison, WI 53706, USA

ARTICLE INFO

Article history:

Received 12 June 2013

Accepted 20 August 2013

ABSTRACT

Using a systematic computer search, nine simple chaotic flows with quadratic nonlinearities were found that have the unusual feature of having a line equilibrium. Such systems belong to a newly introduced category of chaotic systems with hidden attractors that are important and potentially problematic in engineering applications.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

It is widely recognized that mathematically simple systems of nonlinear differential equations can exhibit chaos. With the advent of fast computers, it is now possible to explore the entire parameter space of these systems with the goal of finding parameters that result in some desired characteristics of the system.

Recent research has involved categorizing periodic and chaotic attractors as either self-excited or hidden [1–10]. A self-excited attractor has a basin of attraction that is associated with an unstable equilibrium, whereas a hidden attractor has a basin of attraction that does not intersect with small neighborhoods of any equilibrium points. The classical attractors of Lorenz, Rössler, Chua, Chen, Sprott systems (cases B–S) and other widely-known attractors are those excited from unstable equilibria. From a computational point of view this allows one to use a numerical method in which a trajectory started from a point on the unstable manifold in the neighborhood of an unstable equilibrium, reaches an attractor and identifies it [7]. Hidden attractors cannot be found by this method and are important in engineering applications because they allow unexpected and potentially disastrous responses to perturbations in a structure like a bridge or an airplane wing.

The chaotic attractors in dynamical systems without any equilibrium points or with only stable equilibria are

hidden attractors. That is the reason such systems are rarely found, and only a few such examples have been reported in the literature [11–20].

In this paper, we introduce a new category of chaotic systems with hidden attractors: systems with a line equilibrium. Although in such systems the basin of attraction may intersect the line equilibrium in some sections, there are usually uncountably many points on the line that lie outside the basin of attraction of the chaotic attractor, and thus it is impossible to identify the chaotic attractor for sure by choosing an arbitrary initial condition in the vicinity of the unstable equilibria. In other words, from a computational point of view these attractors are hidden, and knowledge about equilibria does not help in their localization. On the other hand, to the best of our knowledge, although there are dynamical systems with a line equilibrium in the literature [21–24], only one chaotic example has been reported [25], and it is artificial because it is a four-dimensional system that can be reduced to a three-dimensional system in which the line equilibrium vanishes. The goal of this paper is to describe a new category of hidden attractor and expand the list of known mathematically simple hidden chaotic attractors. Thus we perform a systematic computer search for chaos in three-dimensional autonomous systems with quadratic nonlinearities which have been designed so that there will be a line equilibrium, and we ensure that the line equilibrium cannot be made to vanish by reduction to a system of lower dimension.

* Corresponding author.

E-mail address: sajadjafari@aut.ac.ir (S. Jafari).

Table 1
Six simple chaotic flows with line equilibrium.

Case	Equations	(a,b)	Equilibrium	Eigenvalues	LEs	D_{KY}	(x_0, y_0, z_0)
LE ₁	$\dot{x} = y$	$a = 15$	0	$\frac{z \pm \sqrt{z^2 - 4}}{2}$	0.0717	2.1371	0
	$\dot{y} = -x + yz$	$b = 1$	0	0	0		0.5
	$\dot{z} = -x - axy - bxz$		z		-0.5232		0.5
LE ₂	$\dot{x} = y$	$a = 17$	0	$\frac{z \pm \sqrt{z^2 - 4}}{2}$	0.0564	2.1927	0
	$\dot{y} = -x + yz$	$b = 1$	0	0	0		0.4
	$\dot{z} = -y - axy - bxz$		z		-0.2927		0
LE ₃	$\dot{x} = y$	$a = 18$	0	$\frac{z \pm \sqrt{z^2 - 4}}{2}$	0.0556	2.1714	0
	$\dot{y} = -x + yz$	$b = 1$	0	0	0		-0.4
	$\dot{z} = x^2 - axy - bxz$		z		-0.3245		0.5
LE ₄	$\dot{x} = y$	$a = 4$	0	$\frac{z \pm \sqrt{z^2 - 4}}{2}$	0.0539	2.1712	0.2
	$\dot{y} = -x + yz$	$b = 0.6$	0	0	0		0.7
	$\dot{z} = -axy - bxy - yz$		z		-0.3147		0
LE ₅	$\dot{x} = y$	$a = 1.5$	0	$\frac{z \pm \sqrt{z^2 - 4a}}{2}$	0.1386	2.1007	0.7
	$\dot{y} = -ax + yz$	$b = 5$	0	0	0		1
	$\dot{z} = -x^2 - y^2 - bxz$		z		-1.3764		0
LE ₆	$\dot{x} = y$	$a = 0.04$	0	$\frac{z \pm \sqrt{z^2 - 4}}{2}$	0.0543	2.0860	12
	$\dot{y} = -x + yz$	$b = 0.1$	0	0	0		2
	$\dot{z} = -ay^2 - xy - bxz$		z		-0.6314		0
LE ₇	$\dot{x} = z$	$a = 1.85$	0	$\frac{-0.3y \pm \sqrt{0.09y^2 - 4y}}{2}$	0.1144	2.0140	5.1
	$\dot{y} = x + yz$		y	0	0		7
	$\dot{z} = -ax^2 - xy - byz$	$b = 0.3$	0		-1.0270		0
LE ₈	$\dot{x} = z$	$a = 3$	0	$\pm \sqrt{y}$	0.0521	2.0647	0
	$\dot{y} = -x - yz$	$b = 1$	y	0	0		-0.3
	$\dot{z} = ax^2 - xy - bxz$		0		-0.8053		-1
LE ₉	$\dot{x} = z$	$a = 1.62$	x	$\frac{-0.62 \pm \sqrt{6.8644 - 4x^2}}{2}$	0.0642	2.0939	0
	$\dot{y} = -ay + xz$		0	0	0		1
	$\dot{z} = z - bz^2 + xy$	$b = 0.2$	0		-0.6842		0.8




Fig. 1. State space plots of the cases in Table 1 projected onto the xy-plane.

2. Simple chaotic flows with a line equilibrium

In the search for chaotic flows with a line equilibrium, we were inspired by the structure of the conservative Sprott case A system [26],

$$\begin{aligned}\dot{x} &= y \\ \dot{y} &= -x + yz \\ \dot{z} &= 1 - y^2\end{aligned}\quad (1)$$

This system is the oldest and best-known example of a chaotic system with no equilibria, but it does not have a strange attractor since it is conservative. It is an important system since it is a special case of the Nose–Hoover oscillator [27] which describes many natural phenomena [28], and thus it suggests that such systems may have practical as well as theoretical importance.

We consider a general parametric form of Eq. (1) with quadratic nonlinearities of the form

$$\begin{aligned}\dot{x} &= y \\ \dot{y} &= a_1x + a_2yz \\ \dot{z} &= a_3x + a_4y + a_5x^2 + a_6y^2 + a_7xy + a_8xz + a_9yz\end{aligned}\quad (2)$$

As can be seen, this system has a line equilibrium in $(0, 0, z)$ with no other equilibria (in other words the z -axis is the line equilibrium of this system).

An exhaustive computer search was done considering millions of combinations of the coefficients a_1 through a_9 and initial conditions, seeking dissipative cases for which the largest Lyapunov exponent is greater than 0.001. For each case that was found, the space of coefficients was searched for values that are deemed “elegant” [29], by which we mean that as many coefficients as possible are set to zero with the others set to ± 1 if possible or otherwise to a small integer or decimal fraction with the fewest possible digits. Cases LE₁–LE₆ in Table 1 are six simple cases found in this way with only six terms. With some similar procedure, three other similar cases LE₇–LE₉ are included in the table.

In addition to the cases in the table, dozens of additional cases were found, but they were either equivalent to one of the cases listed by some linear transformation of variables, or they were extensions of these cases with more than six terms.

All these cases are dissipative with attractors projected onto the xy -plane as shown in Fig. 1. The equilibria, eigen-




Fig. 2. The largest Lyapunov exponent and bifurcation diagram of case LE₁ showing a period-doubling route to chaos

values, Lyapunov exponent spectra, and Kaplan–Yorke dimensions are shown in Table 1 along with initial conditions that are close to the attractor. As is usual for strange attractors from three-dimensional autonomous systems, the attractor dimension is only slightly greater than 2.0, the largest of which is LE_2 with $D_{KY} = 2.1927$, although no effort was made to tune the parameters for maximum chaos. All the cases appear to approach chaos through a succession of period-doubling limit cycles, a typical example of which (LE_1) is shown in Fig. 2 with decreasing a for $b = 1$. As a decreases further, the strange attractor is destroyed in a boundary crisis.

Fig. 3 shows a cross section in the xz -plane at $y = 0$ of the basin of attraction for the two attractors for the typical case LE_1 . Note that the cross section of the strange attractor nearly touches its basin boundary as is typical of low-dimensional chaotic flows.

For LE_1 we can obtain the eigenvalues from $\lambda(\lambda^2 - z\lambda + 1) = 0$. One of the eigenvalues is zero. The other two depend on z . Using the Routh–Hurwitz stability criterion for the other two, we can say for $z > 0$ the eigenvalues have a positive real part, and thus the positive z -axis is unstable. As shown in Fig. 3, the basin of attraction of the

chaotic attractor intersects the line equilibrium in some portions. However there are other parts of the z -axis (for $z > 0$) which lie in the basin of the stable equilibrium or that attract to infinity. So the strange attractor is hidden in the sense that there are uncountable unstable points on the line equilibrium of which only a tiny portion intersects the basin of the chaotic attractor. In other words, for computational purposes, the attractor is hidden and knowledge about the line equilibrium does not help in its localization.

The reason the stable equilibrium for $z < -2$ does not appear in the basin plot is because it is a node rather than a focus in the xy -plane for $z < -2$. Hence the trajectory never crosses the $y = 0$ plane for $z < -2$. Orbits that start to the left of the equilibrium ($x < -2$) are pulled in the $+z$ direction for $z > -2$ and in the $-z$ direction for $z < -2$. However, they do not go to infinity unless they start far from the equilibrium. Rather, they asymptotically approach it from the $-x$ side and converge to a point on the line that depends on the initial condition. Thus the entire negative z -axis is an attractor, but it is nonlinearly contracting along its length for $-2 < z < 0$ and nonlinearly expanding for $z < -2$.




Fig. 3. Cross section of the basins of attraction of the two attractors in the xz -plane at $y = 0$ for case LE_1 . Initial conditions in the white region lead to unbounded orbits, those in the red region lead to the strange attractor, and those in the light blue region lead to the line equilibrium. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 4. Regions of different dynamic behavior in parameter space for case LE_1 . Light blue represents a static equilibrium, and the black dots correspond to regions of chaos. Each pixel uses a different random condition thereby indicating the coexistence of static and chaotic attractors. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

To show that the behavior described about is not dependent on the particular choice of parameters, Fig. 4 shows the regions of different dynamical behavior in the ab -parameter space. In this plot each pixel represents a different initial condition chosen from a Gaussian distribution with zero mean and unit variance. Thus the region in which the strange attractor (black dots) coexists with the stable line equilibrium (light blue background) extends throughout much of the parameter space. The other eight cases in Table 1 shows similar behavior.

Also of interest is the fact that in cases LE_1 – LE_8 , the strange attractor surrounds the line equilibrium, while in case LE_9 , the line equilibrium lies just outside the strange attractor. In none of the cases does the line equilibrium intersect the attractor, and thus we would not expect homoclinic orbits.

3. Conclusion

In conclusion, it is apparent that simple chaotic systems with a line equilibrium that seemed to be rare, may in fact be rather common. These systems belong to the newly

introduced class of chaotic systems with hidden attractors. In fact they are a new category of them which have not been previously described.

References

- [1] Kuznetsov NV, Leonov GA, Vagitsev VI. Analytical-numerical method for attractor localization of generalized Chua's system. *IFAC Proc.* 2010;4:29–33.
- [2] Kuznetsov NV, Kuznetsova OA, Leonov GA, Vagitsev VI. Hidden attractor in Chua's circuits. In: *ICINCO 2011 – Proc. 8th Int. Conf. Informatics in Control, Automation and Robotics*, 2011. p. 279–283.
- [3] Kuznetsov NV, Leonov GA, Seledzhi SM. Hidden oscillations in nonlinear control systems. *IFAC Proc.* 2011;18:2506–10.
- [4] Leonov GA, Kuznetsov NV. Algorithms for searching for hidden oscillations in the Aizerman and Kalman problems. *Dokl Math* 2011;84:475–81.
- [5] Leonov GA, Kuznetsov NV. Analytical-numerical methods for investigation of hidden oscillations in nonlinear control systems. *IFAC Proc* 2011;18:2494–505.
- [6] Leonov GA, Kuznetsov NV, Kuznetsova OA, Seledzhi SM, Vagitsev VI. Hidden oscillations in dynamical systems. *Trans Syst Control* 2011;6:54–67.
- [7] Leonov GA, Kuznetsov NV, Vagitsev VI. Localization of hidden Chua's attractors. *Phys Lett A* 2011;375:2230–3.
- [8] Leonov GA, Kuznetsov NV, Vagitsev VI. Hidden attractor in smooth Chua systems. *Physica D* 2012;241:1482–6.

- [9] Leonov GA, Kuznetsov NV. Analytical-numerical methods for hidden attractors' localization: the 16th Hilbert problem, Aizerman and Kalman conjectures, and Chua circuits. In: *Numerical Methods for Differential Equations, Optimization, and Technological Problems, Computational Methods in Applied Sciences*, vol. 27. 2013. p. 41–64.
- [10] Leonov GA, Kuznetsov NV. Hidden attractors in dynamical systems: from hidden oscillation in Hilbert–Kolmogorov, Aizerman and Kalman problems to hidden chaotic attractor in Chua circuits. *Int. J. Bifurcation Chaos* 2013;23:1330002. p. 69.
- [11] Jafari S, Sprott JC, Golpayegani SMRH. Elementary quadratic chaotic flows with no equilibria. *Phys Lett A* 2013;377:699–702.
- [12] Molaie M, Jafari S, Sprott JC, Golpayegani SMRH. Simple chaotic flows with one stable equilibrium. *Int. J. Bifurcation Chaos*, in press.
- [13] Wang X, Chen G. A chaotic system with only one stable equilibrium. *Commun Nonlinear Sci Numer Simulat* 2012;17:1264–72.
- [14] Wang X, Chen G. Constructing a chaotic system with any number of equilibria. *Nonlinear Dyn* 2013;71:429–36.
- [15] Wei Z, Yang Q. Dynamical analysis of the generalized Sprott C system with only two stable equilibria. *Nonlinear Dyn* 2012;68:543–54.
- [16] Wang X, Chen J, Lu JA, Chen G. A simple yet complex one-parameter family of generalized Lorenz-like systems. *Int J Bifurcation Chaos* 2012;22:1250116. p. 16.
- [17] Wei Z. Dynamical behaviors of a chaotic system with no equilibria. *Phys Lett A* 2011;376:102–8.
- [18] Wang Z, Cang S, Ochola EO, Sun Y. A hyperchaotic system without equilibrium. *Nonlinear Dyn* 2012;69:531–7.
- [19] Wei Z. Delayed feedback on the 3-D chaotic system only with two stable node-foci. *Comput Math Appl* 2011;63:728–38.
- [20] Wei Z, Yang Q. Anti-control of Hopf bifurcation in the new chaotic system with two stable node-foci. *Appl Math Comput* 2010;217:422–9.
- [21] Messias M, Nespole C, Botta VA. Hopf bifurcation from lines of equilibria without parameters in memristor oscillators. *Int. J. Bifurcation Chaos* 2010;20:437–50.
- [22] Fiedler B, Liebscher S, Alexander JC. Generic hopf bifurcation from lines of equilibria without parameters: I theory. *J Differ Equ* 2000;167:16–35.
- [23] Fiedler B, Liebscher S. Generic hopf bifurcation from lines of equilibria without parameters: II. Systems of viscous hyperbolic balance laws. *SIAM J Math Anal* 1998;31:1396–404.
- [24] Fiedler B, Liebscher S, Alexander JC. Generic hopf bifurcation from lines of equilibria without parameters III: binary oscillations. *Int J Bifurcation Chaos* 2000;10:1613.
- [25] Zhou P, Huang K, Yang CD. A fractional-order chaotic system with an infinite number of equilibrium points. *Discrete Dyn Nat Soc* 2013;6. Article ID 910189.
- [26] Sprott JC. Some simple chaotic flows. *Phys Rev E* 1994;50:R647.
- [27] Hoover WG. Remark on some simple chaotic flows. *Phys Rev E* 1995;51:759.
- [28] Posh HA, Hoover WG, Vesely FJ. Canonical dynamics of the nosé oscillator: stability, order, and chaos. *Phys Rev A* 1986;33:4253.
- [29] Sprott JC. *Elegant chaos: algebraically simple chaotic flows*. World Scientific; 2010.